

## 鸟嘌呤解离刺激因子样蛋白 1 抗体

产品货号： mIR19847

英文名称： RGL1

中文名称： 鸟嘌呤解离刺激因子样蛋白 1 抗体

别名： KIAA0959; Ral guanine nucleotide dissociation stimulator like 1; RalGDS like 1; RalGDS like; RGL 1; RGL. RGL1\_HUMAN

研究领域： 肿瘤 细胞生物 信号转导 G 蛋白信号

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Pig, Cow, Horse, Sheep,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 87kDa

细胞定位： 细胞浆

性状： Lyophilized or Liquid

浓度： 1mg/ml

**免 疫 原：** KLH conjugated synthetic peptide derived from human RGL1:601-676/768

**亚 型：** IgG

**纯化方法：** affinity purified by Protein A

**储 存 液：** 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

**保存条件：** Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

**PubMed：** PubMed

**产品介绍：** RGL1 (Ral guanine nucleotide dissociation stimulator-like 1) is a putative guanine nucleotide exchange factor involved in Ras and Ral signaling pathways as a downstream effector protein.

**Function:**

Probable guanine nucleotide exchange factor.

**Subunit:**

Interacts with Ras.

**Tissue Specificity:**

Expressed in a wide variety of tissues with strong expression being seen in the heart, brain, kidney, spleen and testis.

**Similarity:**

Contains 1 N-terminal Ras-GEF domain.

Contains 1 Ras-associating domain.

Contains 1 Ras-GEF domain.

**SWISS:**

Q9NZL6

**Gene ID:**

23179

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.