

鸟嘌呤解离刺激因子样蛋白 2 抗体

产品货号： mlR19848

英文名称： RGL2

中文名称： 鸟嘌呤解离刺激因子样蛋白 2 抗体

别名： GDS related protein; HKE1.5; KE1.5; RAB2L; Ral guanine nucleotide dissociation stimulator-like 2; RalGDS-like 2; RalGDS-like factor; Ras-associated protein RAB2L; RGL2; RGL2_HUMAN.

研究领域： 细胞生物 信号转导 G 蛋白信号

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat,

产品应用： WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500

(石蜡切片需做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 84kDa

细胞定位： 细胞核

性状： Lyophilized or Liquid

浓度： 1mg/ml

免疫原： KLH conjugated synthetic peptide derived from human RGL2:351-450/777

亚型： IgG

纯化方法： affinity purified by Protein A

储存液： 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件： Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed： PubMed