

锌指蛋白 165 抗体

产品货号： mlR18503

英文名称： ZNF165

中文名称： 锌指蛋白 165 抗体

别 名： Cancer/testis antigen 53; CT53; LD65; MGC140342; OTTHUMP00000016195; Zinc finger and SCAN domain containing protein 7; Zinc finger and SCAN domain-containing protein 7; Zinc finger protein 165; ZINC FINGER PROTEIN 165; ZNF165; ZN165_HUMAN; ZNF165; ZPF165; ZSCAN7.

研究领域： 转录调节因子 锌指蛋白 表观遗传学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量： 56kDa

细胞定位： 细胞核

性 状： Lyophilized or Liquid

浓 度： 1mg/ml

免 疫 原： KLH conjugated synthetic peptide derived from human ZNF165:1-100/485

亚 型： IgG

纯化方法： affinity purified by Protein A

储 存 液： 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件： Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed： PubMed

产品介绍： This gene encodes a member of the Kruppel family of zinc finger proteins. Members of this DNA-binding protein family act as transcriptional regulators. This gene is located within a cluster of zinc finger family members. The encoded protein may play a role in spermatogenesis. [provided by RefSeq, Jul 2008]

Function:

May be involved in transcriptional regulation.

Subcellular Location:

Nucleus.

Tissue Specificity:

Expressed specifically in testis.

Similarity:

Belongs to the krueppel C2H2-type zinc-finger protein family.

Contains 6 C2H2-type zinc fingers.

Contains 1 SCAN box domain.

SWISS:

P49910

Gene ID:

7718

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.