

## 锌指蛋白 213 抗体

产品货号: mlR12800

英文名称: ZNF213

中文名称: 锌指蛋白 213 抗体

别 名: CR53; Putative transcription factor CR53; zinc finger protein 213; Zinc finger protein with KRAB and SCAN domains 21; ZKSCAN21.

研究领域: 细胞生物 转录调节因子 锌指蛋白 表观遗传学

抗体来源: Rabbit

克隆类型: Polyclonal

交叉反应: Human, Mouse, Rat, Cow, Rabbit,

**产品应用 :** ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 (石蜡切片需

做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量: 51kDa

细胞定位: 细胞核

性 状: Lyophilized or Liquid

浓 度: 1mg/ml



Belongs to the krueppel C2H2-type zinc-finger protein family.

Contains 5 C2H2-type zinc fingers.

免疫原: KLH conjugated synthetic peptide derived from human ZNF213:301-400/459
亚 型: IgG
纯化方法: affinity purified by Protein A
储 存 液: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.
保存条件: Store at -20 $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20 $^{\circ}$ C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 $^{\circ}$ C.
PubMed: PubMed
产品介绍: C2H2 zinc finger proteins, such as ZNF213, have bipartite structures in which one domain binds DNA or RNA and the other modulates target gene expression.[supplied by OMIM, Apr 2004]
Function:
May be involved in transcriptional regulation.
Subcellular Location:
Nuclear.
Tissue Specificity:
Widely expressed with highest levels in testis.
Similarity:



applications.

Contains 1 KRAB domain.
Contains 1 SCAN box domain.
SWISS:
O14771
Gene ID:
7760
Important Note:
This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic