

p53 调节凋亡诱导蛋白 1 抗体

产品货号： mlR7094

英文名称： p53AIP1

中文名称： p53 调节凋亡诱导蛋白 1 抗体

别 名： Alternative names p53 regulated apoptosis inducing protein 1; p53-regulated apoptosis-inducing protein 1; P53AIP 1; p53 AIP1; TP53AIP 1; TP53AIP1; TPIP1_HUMAN.

研究领域： 肿瘤 细胞生物 细胞凋亡

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse,

产品应用： WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量：13.6kDa

细胞定位：细胞浆

性状：Lyophilized or Liquid

浓度：1mg/ml

免疫原：KLH conjugated synthetic peptide derived from human p53AIP1:65-124/124

亚型：IgG

纯化方法：affinity purified by Protein A

储存液：0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件：Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PubMed：PubMed

产品介绍：This gene is specifically expressed in the thymus, and encodes a protein that is localized to the mitochondrion. The expression of this gene is inducible by p53, and it is thought to play an important role in

mediating p53-dependent apoptosis. Alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Oct 2011].

Function:

May play an important role in mediating p53/TP53-dependent apoptosis.

Subcellular Location:

Mitochondrion.

Tissue Specificity:

Only found to be expressed in thymus.

SWISS:

Q9HCN2

Gene ID:

63970

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

产品图片：

