

p53 和 DNA 损伤调节蛋白 1 抗体

产品货号： mlR12604

英文名称： PDRG1

中文名称： p53 和 DNA 损伤调节蛋白 1 抗体

别名： C20orf126; p53 and DNA damage regulated 1; p53 and DNA damage regulated protein 1; PDRG.

研究领域： 细胞生物 信号转导 表观遗传学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Dog, Horse, Rabbit,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 15kDa

细胞定位： 细胞浆

性状： Lyophilized or Liquid

浓度： 1mg/ml

免疫原： KLH conjugated synthetic peptide derived from human PDRG1:1-100/133

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed : PubMed

产品介绍 : May play a role in chaperone-mediated protein folding. Predominantly expressed in testis.

Function:

May play a role in chaperone-mediated protein folding

Subcellular Location:

Cytoplasmic.

Tissue Specificity:

Predominantly expressed in normal testis and exhibits reduced but detectable expression in other organs.

Similarity:

Belongs to the prefoldin subunit beta family.

SWISS:

Q9NUG6

Gene ID:

81572

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.