



## OSTC 蛋白抗体

产品货号 : mlR17530

英文名称 : OSTC

中文名称 : OSTC 蛋白抗体

别 名 : DC2; HDCMD45P; HSPC307; Hydrophobic protein HSF-28; Oligosaccharyltransferase complex subunit; Oligosaccharyltransferase complex subunit OSTC; OSTC; OSTC\_HUMAN.

研究领域 : 细胞生物 信号转导 新陈代谢

抗体来源 : Rabbit

克隆类型 : Polyclonal

交叉反应 : Human, Mouse, Rat, Dog, Pig, Horse, Rabbit, Zebrafish, Xenopus laevis

产品应用 : ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 (石蜡切片需做抗原修复)



not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 17kDa

细胞定位： 细胞膜

性 状： Lyophilized or Liquid

浓 度： 1mg/ml

免 疫 原： KLH conjugated synthetic peptide derived from human OSTC:21-100/149 <Extracellular>

亚 型： IgG

纯化方法： affinity purified by Protein A

储 存 液： 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件： Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.



**PubMed :** PubMed

**产品介绍 :** Belongs to the OSTC family.

**Subcellular Location:**

Membrane.

**SWISS:**

Q9NRP0

**Gene ID:**

58505

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.