



甲基 CpG 结合域蛋白 3 样 2 抗体

产品货号 : mlR18702

英文名称 : MBD3L2

中文名称 : 甲基 CpG 结合域蛋白 3 样 2 抗体

别 名 : MB3L2_HUMAN; MBD3 like 2; MBD3-like protein 2; Mbd3l2; Methyl CpG binding domain protein 3 like 2; Methyl-CpG-binding domain protein 3-like 2.

研究领域 : 转录调节因子 结合蛋白 表观遗传学

抗体来源 : Rabbit

克隆类型 : Polyclonal

交叉反应 : Human,

产品应用 : ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 (石蜡切片需做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量 : 23kDa

细胞定位 : 细胞核

性 状 : Lyophilized or Liquid

浓 度 : 1mg/ml



免 疫 原 : KLH conjugated synthetic peptide derived from human MBD3L2:131-208/208

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed : PubMed

产品介绍 : This gene encodes a protein that is related to methyl-CpG-binding proteins but lacks the methyl-CpG binding domain. The protein has been found in germ cell tumors and some somatic tissues. [provided by RefSeq, Jul 2008]

Tissue Specificity:

Detected at low levels in several somatic tissues. Highly expressed in the ovarian teratocarcinoma cell line PA-1.

Similarity:

Belongs to the MBD3L family.

SWISS:

Q8NHZ7

Gene ID:

125997



Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.