

## 基质细胞衍生因子 2 样蛋白 1 抗体

产品货号： mlR4446

英文名称： SDF2L1

中文名称： 基质细胞衍生因子 2 样蛋白 1 抗体

别名： PWP1-interacting protein 8; SDF2-like protein 1; SDF2L\_HUMAN; SDF2L1; Stromal cell-derived factor 2-like protein 1.

研究领域： 细胞生物 信号转导 转运蛋白

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Dog, Cow, Sheep,

产品应用： WB=1:500-2000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量 : 24kDa

细胞定位 : 细胞浆

性 状 : Lyophilized or Liquid

浓 度 : 1mg/ml

免 疫 原 : KLH conjugated synthetic peptide derived from human SDF2L1:121-221/221

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : Preservative: 15mM Sodium Azide, Constituents: 1% BSA, 0.01M PBS, pH 7.4

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed : PubMed

**产品介绍 background:**

SDF2L1 (Stromal Cell Derived Factor 2 Like 1) is a Protein Coding gene. GO annotations related to this gene include chaperone binding and misfolded protein binding. An important paralog of this gene is SDF2.

**Subcellular Location:**

Endoplasmic reticulum lumen.

**Tissue Specificity:**

Ubiquitously expressed with high expression in testis, moderate expression in the pancreas, spleen, prostate, small intestine and colon. Very low expression is seen in brain and skeletal muscle.

**Similarity:**

Contains 3 MIR domains.

**SWISS:**

Q9HCN8

**Gene ID:**

23753

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

产品图片

