

三磷酸腺苷门控阳离子通道蛋白抗体

产品货号： mlR7690

英文名称： P2RX4

中文名称： 三磷酸腺苷门控阳离子通道蛋白抗体

别名： ATP gated cation channel protein; ATP receptor; p2rx4; P2RX4_HUMAN; P2X purinoceptor 4; P2X receptor subunit 4; P2X4; P2X4R; Purinergic receptor; Purinergic receptor P2X ligand gated ion channel 4; Purinergic receptor P2X4; Purinoceptor P2X4.

研究领域： 肿瘤 心血管 细胞生物 神经生物学 信号转导 通道蛋白

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Dog, Cow,

产品应用： WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量 : 43kDa

细胞定位 : 细胞膜

性 状 : Lyophilized or Liquid

浓 度 : 1mg/ml

免 疫 原 : KLH conjugated synthetic peptide derived from human P2RX4:238-338/388 <Extracellular>

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PubMed : PubMed

产品介绍： The P2X receptor family is comprised of ligand-gated ion channels that allow for the increased permeability of calcium into the cell in response to extracellular ATP. The seven P2X receptors, P2X1-P2X7, form either homomeric or heteromeric channels or both. They are characterized by intracellular amino- and carboxy-termini. P2X receptors are expressed in a wide variety of tissues, including neurons, prostate, bladder, pancreas, colon, testis and ovary. The major function of the P2X receptors is to mediate synaptic transmissions between neurons and to other tissues via the binding of extracellular ATP, which acts as a neurotransmitter. The P2X receptors may be involved in the onset of necrosis or apoptosis after prolonged exposure to high concentrations of extracellular ATP.

Function:

Receptor for ATP that acts as a ligand-gated ion channel. This receptor is insensitive to the antagonists PPADS and suramin.

Subunit:

Functional P2XRs are organized as homomeric and heteromeric trimers.

Subcellular Location:

Membrane; Multi-pass membrane protein.

Similarity:

Belongs to the P2X receptor family.

SWISS:

Q99571

Gene ID:

5025

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

产品图片：

