

NOTO 蛋白抗体

产品货号： mIR19321

英文名称： NOTO

中文名称： NOTO 蛋白抗体

别名： floating head homolog; Homeobox protein notochord; Not; Not homeodomain protein; Noto; NOTO_HUMAN; notochord homolog (Xenopus laevis).

研究领域： 发育生物学 神经生物学 转录调节因子 表观遗传学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Pig, Rabbit,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 27kDa

细胞定位： 细胞核

性状： Lyophilized or Liquid

浓度： 1mg/ml

免疫原： KLH conjugated synthetic peptide derived from human NOTO:121-220/251

亚型： IgG

纯化方法： affinity purified by Protein A

储 存 液： 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件： Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20 ° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed： PubMed

产品介绍： Transcription regulator acting downstream of both FOXA2 and T during notochord development. Required for node morphogenesis. Is essential for cilia formation in the posterior notochord (PNC) and for left-right patterning; acts upstream of FOXJ1 and RFX3 in this process and is required for the expression of various components important for axonemal assembly and function. Plays a role in regulating axial versus paraxial cell fate.

Subcellular Location:

Nucleus.

Similarity:

Contains 1 homeobox DNA-binding domain.

Gene ID:

344022

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.