

NMDA 受体调节 1 样蛋白抗体

产品货号： mIR19022

英文名称： NARG1L

中文名称： NMDA 受体调节 1 样蛋白抗体

别 名： DKFZp686O08147; MGC40612; N alpha acetyltransferase 16, NatA auxiliary subunit; N(alpha) acetyltransferase 16, NatA auxiliary subunit; NARG1 like protein; NAT2; NMDA receptor regulated 1 like; NMDA receptor regulated 1 like protein; OTTHUMP00000018318; OTTHUMP00000018320; OTTHUMP000000198977; RP11-396A22.1.

研究领域： 细胞生物 免疫学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Chicken, Dog, Pig, Cow, Horse, Rabbit,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量： 101kDa

细胞定位： 细胞浆

性 状： Lyophilized or Liquid

浓 度 : 1mg/ml

免 疫 原 : KLH conjugated synthetic peptide derived from human NARG1L:551-650/864

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed : PubMed

产品介绍 : The NARG1-like protein (NARG1L) is an 864 amino acid protein that contains 7 TPR repeats and may also be a component of a complex that displays N-terminal acetyltransferase activity.

Function:

Auxillary subunit of the N-terminal acetyltransferase A (Nata) complex which displays alpha (N-terminal) acetyltransferase activity.

Subunit:

Component of the N-terminal acetyltransferase A (NatA) complex composed of NAA10 and NAA16.

Similarity:

Contains 7 TPR repeats.

SWISS:

Q6N069

Gene ID:

79612

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.