

## 甲状腺激素受体相关蛋白复合物 230 样蛋白抗体

产品货号： mlR18792

英文名称： MED12L

中文名称： 甲状腺激素受体相关蛋白复合物 230 样蛋白抗体

别名： MD12L\_HUMAN; MED12L; Mediator complex subunit 12-like protein; Mediator of RNA polymerase II transcription subunit 12-like protein; NOPAR; Thyroid hormone receptor-associated-like protein; TNRC11L; TRALP; TRALPUSH; Trinucleotide repeat-containing gene 11 protein-like.

研究领域： 细胞生物 转录调节因子 表观遗传学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 240kDa

细胞定位： 细胞核

性状： Lyophilized or Liquid

浓 度 : 1mg/ml

免 疫 原 : KLH conjugated synthetic peptide derived from human MED12L:601-700/2145

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed : PubMed

产品介绍 : The protein encoded by this gene is part of the Mediator complex, which is involved in transcriptional coactivation of nearly all RNA polymerase II-dependent genes. The Mediator complex links gene-specific transcriptional activators with the basal transcription machinery. [provided by RefSeq, May 2010]

**Function:**

May be a component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors.

**Subunit:**

Belongs to the Mediator complex subunit 12 family.

**Subcellular Location:**

Nucleus.

**Similarity:**

Belongs to the Mediator complex subunit 12 family.

**SWISS:**

Q86YW9

**Gene ID:**

116931

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.