

胰肠同源结构域转录因子抗体

产品货号: mIR18172

英文名称: ISX/Intestine specific homeobox

中文名称: 胰肠同源结构域转录因子抗体

别 名: DKFZp781N2395; Intestine-specific homeobox; Isx; ISX_HUMAN; MGC138417; OTTHUMP00000028558; Pancreas intestine homeodomain transcription factor; Pix 1; RAX like homeobox; RAX-like homeobox; RAXLX.

研究领域: 细胞生物 转录调节因子 转运蛋白 表观遗传学

抗体来源: Rabbit

克隆类型: Polyclonal

交叉反应: Human, Mouse, Rat, Cow, Horse, Sheep,

产品应用 : WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500

(石蜡切片需做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量: 27kDa

细胞定位: 细胞核

性 状: Lyophilized or Liquid

浓 度: 1mg/ml

免疫原: KLH conjugated synthetic peptide derived from human ISX/Intestine specific homeobox:81-

180/245



亚 型: IgG

纯化方法: affinity purified by Protein A

储存液: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件: Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20 °C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PubMed: PubMed

产品介绍: Homeobox genes encode DNA-binding proteins, many of which are thought to be involved in early embryonic development. Homeobox genes encode a DNA-binding domain of 60 to 63 amino acids referred to as the homeodomain. This gene is a member of the RAXLX homeobox gene family. [provided by RefSeq, Jul 2008]

Function:

Transcription factor that regulates gene expression in intestine. May participate in vitamin A metabolism most likely by regulating BCMO1 expression in the intestine.

Subcellular Location:

Nucleus.

Similarity:

Contains 1 homeobox DNA-binding domain.

SWISS:

Q2M1V0



91464

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

产品图片

