

## NDUFC2 蛋白抗体

产品货号： mIR19090

英文名称： NDUFC2

中文名称： NDUFC2 蛋白抗体

别名： B14.5b; CI-B14.5b; Complex I B14.5b; Complex I subunit B14.5b; Complex I-B14.5b; HLC-1; HLC1; Human lung cancer oncogene 1 protein; NADH dehydrogenase (ubiquinone) 1, subcomplex unknown, 2, 14.5kDa; NADH dehydrogenase [ubiquinone] 1 subunit C2; NADH dehydrogenase ubiquinone 1 subunit C2; NADH ubiquinone oxidoreductase subunit B14.5b; NADH-ubiquinone oxidoreductase subunit B14.5b; NADHDH2; NDUC2\_HUMAN; NDUFC2.

研究领域： 肿瘤 细胞生物 信号转导

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Mouse, Rat, Cow, Horse, Rabbit, Sheep,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 14kDa

细胞定位： 细胞浆

性状： Lyophilized or Liquid

浓 度 : 1mg/ml

免 疫 原 : KLH conjugated synthetic peptide derived from human NDUFC2:1-80/119

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed : PubMed