

## 睾丸分化过程转录因子 MRO 抗体

产品货号： mIR17764

英文名称： MRO

中文名称： 睾丸分化过程转录因子 MRO 抗体

别名： B29; beside the Ma29 deletion; C18orf3; Male-specific transcription in the developing reproductive organs; Mro; MSTRO\_HUMAN; Protein B29; Protein maestro.

研究领域： 细胞生物 发育生物学 转录调节因子 细胞分化

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量 : 29kDa

细胞定位 : 细胞核

性 状 : Lyophilized or Liquid

浓 度 : 1mg/ml

免 疫 原 : KLH conjugated synthetic peptide derived from human MRO:1-100/248

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

**PubMed :** PubMed

**产品介绍 :** This gene is specifically transcribed in males before and after differentiation of testis, and the encoded protein may play an important role in a mammalian sex determination. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]

**Subcellular Location:**

Nucleus > nucleolus.

**Tissue Specificity:**

Ubiquitous.

**Similarity:**

Contains 1 HEAT repeat.

**SWISS:**

Q9BYG7

**Gene ID:**

83876

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

