



MYCNOS 蛋白抗体

产品货号 : mlR19141

英文名称 : MYCNOS

中文名称 : MYCNOS 蛋白抗体

别 名 : MYCN opposite strand/antisense RNA (non-protein coding); MYCN-AS1; N-cym protein; NCYM; NYCM.

研究领域 : 肿瘤 细胞生物 免疫学 神经生物学

抗体来源 : Rabbit

克隆类型 : Polyclonal

交叉反应 : Human,

产品应用 : ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 (石蜡切片需做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量 : 12kDa

细胞定位 : 细胞浆

性 状 : Lyophilized or Liquid

浓 度 : 1mg/ml



免 疫 原 : KLH conjugated synthetic peptide derived from human MYCNOS:31-109/109

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed : PubMed

产品介绍 : This gene is located on the opposite strand of v-myc avian myelocytomatosis viral oncogene neuroblastoma derived homolog (GeneID 4613). Transcription of this antisense gene may influence the progression of neuroblastoma by inhibiting transcription of v-myc avian myelocytomatosis viral oncogene neuroblastoma derived homolog. This gene is represented as non-coding pending further experimental evidence of a protein product. Alternatively spliced transcript variants have been described. [provided by RefSeq, Jan 2014]

Function:

May have a functional role during normal fetal development.

Tissue Specificity:

Expressed in the fetal brain, lung, liver and kidney at varying low levels.

SWISS:

P40205



Gene ID:

10408

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.