



粒细胞趋化蛋白 2/GCP2 抗体

产品货号 : mlR1793

英文名称 : CXCL6/GCP2

中文名称 : 粒细胞趋化蛋白 2/GCP2 抗体

别 名 : granulocyte chemotactic protein 2; C X C motif chemokine 6; Chemokine (C X C motif) ligand 6; Chemokine alpha 3; CKA 3; CKA3; CXCL 6; CXCL6; GCP 2; GCP2; GCP-2; Granulocyte chemotactic protein 2; SCYB6; Small inducible cytokine B6; Small inducible cytokine subfamily B (Cys X Cys) member 6; Small inducible cytokine subfamily B (Cys X Cys) member b.

研究领域 : 细胞生物 免疫学 趋化因子

抗体来源 : Rabbit

克隆类型 : Polyclonal

交叉反应 : Human, Cow, Horse,

产品应用 : WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 IF=1:100-500 (石蜡切片需做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量 : 12kDa

细胞定位 : 分泌型蛋白

性 状 : Lyophilized or Liquid

浓 度 : 1mg/ml

免 疫 原 : KLH conjugated synthetic peptide derived from human CXCL6:51-114/114



亚型 : IgG

纯化方法 : affinity purified by Protein A

储存液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed : PubMed

产品介绍 : Neutrophil recruitment to inflammatory sites is mediated by two related receptors: CXCR1 and CXCR2. Both receptors share two ligands, interleukin-8 (CXCL8) and GCP2 (CXCL6). GCP2 belongs to the intercrine alpha (chemokine CxC) family.

Function:

Chemotactic for neutrophil granulocytes.

Subcellular Location:

Secreted.

Similarity:

Belongs to the intercrine alpha (chemokine CxC) family.

SWISS:

P50228



Gene ID:

6372

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.