



抑癌基因 PDLIM4 抗体

产品货号 : mlR6093

英文名称 : RIL

中文名称 : 抑癌基因 PDLIM4 抗体

别 名 : Enigma homolog; LIM domain protein; LIM protein RIL; PDLIM 4; PDLIM4; PDZ and LIM domain 4; PDZ and LIM domain protein 4; Reversion induced LIM protein; PDLI4_HUMAN.

研究领域 : 肿瘤 染色质和核信号 转录调节因子

抗体来源 : Rabbit

克隆类型 : Polyclonal

交叉反应 : Human, Mouse, Rat, Chicken, Dog, Pig, Cow, Horse, Rabbit,

产品应用 : ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 IF=1:100-500 (石蜡切片需做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.



分子量： 35kDa

细胞定位： 细胞浆

性 状： Lyophilized or Liquid

浓 度： 1mg/ml

免 疫 原： KLH conjugated synthetic peptide derived from human RIL:251-330/330

亚 型： IgG

纯化方法： affinity purified by Protein A

储 存 液： 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件： Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PubMed： PubMed

产品介绍 background:



RIL, also known as PDLIM4 is a putative tumor suppressor gene. RIL is transcriptionally repressed by hypermethylation in a variety of tumors including breast and prostate. The difference in methylation status of RIL between normal and cancerous tissues is being investigated as to determine the use of RIL as a potential biomarker. Two isoforms of human RIL one have been identified.

Subcellular Location:

Cytoplasmic

Similarity:

Contains 1 LIM zinc-binding domain.

Contains 1 PDZ (DHR) domain.

SWISS:

P50479

Gene ID:

8572

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

产品图片：

