

## 精子发生相关蛋白 20 抗体

产品货号： mlR17625

英文名称： SPATA20

中文名称： 精子发生相关蛋白 20 抗体

别名： SPATA20; Sperm protein SSP411; Sperm-specific protein 411; Spermatogenesis associated 20; Spermatogenesis-associated protein 20; SPT20\_HUMAN; Ssp411; Tisp78; Transcript increased in spermiogenesis 78.

研究领域： 细胞生物 发育生物学 干细胞

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Dog, Pig, Cow, Horse, Sheep, Guinea Pig,

产品应用： WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500

(石蜡切片需做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量 : 58kDa

细胞定位 : 细胞外基质 分泌型蛋白

性 状 : Lyophilized or Liquid

浓 度 : 1mg/ml

免 疫 原 : KLH conjugated synthetic peptide derived from human SPATA20:431-530/786

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

**PubMed :** PubMed

**产品介绍 :** SPATA20 is a 786 amino acid protein that may be involved in fertility regulation. SPATA20 is expressed in testis in an age dependent manner and localizes to round and elongated spermatids. SPATA20 contains a conserved thioredoxin-like domain near the N-terminus region and exists as four isoforms produced by alternative splicing events. SPATA20 is encoded by a gene located on human chromosome 17, which comprises over 2.5% of the human genome and encodes over 1,200 genes. Two key tumor suppressor genes are associated with chromosome 17, namely, p53 and BRCA1. Tumor suppressor p53 is necessary for maintenance of cellular genetic integrity by moderating cell fate through DNA repair versus cell death. Malfunction or loss of p53 expression is associated with malignant cell growth and Li-Fraumeni syndrome. Like p53, BRCA1 is directly involved in DNA repair, though specifically it is recognized as a genetic determinant of early onset breast cancer and predisposition to cancers of the ovary, colon, prostate gland and fallopian tubes.

**Function:**

May play a role in fertility regulation.

**Subcellular Location:**

Secreted.

**SWISS:**

Q8TB22

**Gene ID:**

64847

**Important Note:**



This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.