

## 磷酸化谷氨酸受体 2B 抗体

产品货号： mlR3306

英文名称： Phospho-NMDAR2B (Tyr1070)

中文名称： 磷酸化谷氨酸受体 2B 抗体

别名： NMDAR2B (phospho Y1070); p-NMDAR2B (phospho Y1070); NMDAR2B (Phospho-Tyr1070); NMDAR2B (Phospho-Y1070); p-NMDAR2B (Tyr1070); p-NMDAR2B (Y1070); AW490526; Glutamate [NMDA] receptor subunit epsilon 2; Glutamate [NMDA] receptor subunit epsilon-2; Glutamate Receptor Ionotropic N Methyl D Aspartate 2B; Glutamate Receptor Ionotropic N Methyl D Aspartate subunit 2B; Glutamate receptor ionotropic NMDA2B; Glutamate receptor subunit epsilon 2; Glutamate receptor, ionotropic, NMDA2B (epsilon 2); GRIN 2B; GRIN2B; hNR 3; hNR3; MGC142178; MGC142180; N methyl D aspartate receptor channel subunit epsilon 2; N METHYL D ASPARTATE RECEPTOR CHANNEL SUBUNIT EPSILON 2; N methyl D aspartate receptor subtype 2B; N methyl D aspartate receptor subunit 2B; N methyl D aspartate receptor subunit 3; N-methyl D-aspartate receptor subtype 2B; N-methyl-D-aspartate receptor subunit 3; NMDA NR2B; NMDA R2B; Nmdar2b; NMDE2; NMDE2\_HUMAN; NME2; NR2B; NR3.

产品类型： 磷酸化抗体

研究领域： 细胞生物 免疫学 神经生物学 信号转导 细胞膜受体

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Chicken, Dog, Cow, Horse,

产品应用： WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量：164kDa

细胞定位：细胞膜

性状：Lyophilized or Liquid

浓度：1mg/ml

免疫原：KLH conjugated Synthesised phosphopeptide derived from human NMDAR2B around the phosphorylation site of Tyr1070:VT(p-Y)GN

亚型：IgG

纯化方法：affinity purified by Protein A

储存液：0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件：Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20 ° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

**PubMed：** PubMed

**产品介绍：** N-methyl-D-aspartate (NMDA) receptors are a class of ionotropic glutamate receptors. NMDA receptor channel has been shown to be involved in long-term potentiation, an activity-dependent increase in the efficiency of synaptic transmission thought to underlie certain kinds of memory and learning. NMDA receptor channels are heteromers composed of three different subunits: NR1 (GRIN1), NR2 (GRIN2A, GRIN2B, GRIN2C, or GRIN2D) and NR3 (GRIN3A or GRIN3B). The NR2 subunit acts as the agonist binding site for glutamate. This receptor is the predominant excitatory neurotransmitter receptor in the mammalian brain. [provided by RefSeq, Jul 2008].

**Function:**

NMDA receptor subtype of glutamate-gated ion channels with high calcium permeability and voltage-dependent sensitivity to magnesium. Mediated by glycine. In concert with DAPK1 at extrasynaptic sites, acts as a central

mediator for stroke damage. Its phosphorylation at Ser-1303 by DAPK1 enhances synaptic NMDA receptor channel activity inducing injurious  $\text{Ca}^{2+}$  influx through them, resulting in an irreversible neuronal death (By similarity).

**Subunit:**

Forms heteromeric channel of a zeta subunit (GRIN1), a epsilon subunit (GRIN2A, GRIN2B, GRIN2C or GRIN2D) and a third subunit (GRIN3A or GRIN3B). Found in a complex with GRIN1 and GRIN3B. Found in a complex with GRIN1, GRIN3A and PPP2CB. Interacts with PDZ domains of INADL and DLG4. Interacts with HIP1 and NETO1. Interacts with MAGI3. Interacts with DAPK1.

**Subcellular Location:**

Cell membrane; Multi-pass membrane protein. Cell junction, synapse, postsynaptic cell membrane; Multi-pass membrane protein.

**Tissue Specificity:**

Primarily found in the fronto-parieto-temporal cortex and hippocampus pyramidal cells, lower expression in the basal ganglia.

**Post-translational modifications:**

Phosphorylation at Ser-1303 by DAPK1 enhances synaptic NMDA receptor channel activity.

**DISEASE:**

Defects in GRIN2B are the cause of mental retardation autosomal dominant type 6 (MRD6) [MIM:613970]. Mental retardation is characterized by significantly below average general intellectual functioning associated with impairments in adaptive behavior and manifested during the developmental period. Note=Chromosomal aberrations involving GRIN2B have been found in patients with mental retardation. Translocations t(9;12)(p23;p13.1) and t(10;12)(q21.1;p13.1) with a common breakpoint in 12p13.1.

**Similarity:**

Belongs to the glutamate-gated ion channel (TC 1.A.10.1) family. NR2B/GRIN2B subfamily.

**SWISS:**

Q13224

**Gene ID:**

2904

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

谷氨酸受体（NR2B）是脊椎动物中枢神经系统兴奋型神经传递的主要介质。在突触可塑性极大脑学习及记忆功能方面起关键作用。