

RhoB 蛋白抗体

产品货号： mlR11142

英文名称： Rho B

中文名称： RhoB 蛋白抗体

别名： Rh -related GTP binding protein RhoB; Rh -related GTP binding protein RhoC; RHOH6; Transforming protein RhoA; AA017882 antibody Aplysia ras related homolog 9 (RhoC); ARH9; ARHA; ARHB; ARHC; H6; MST081; MSTP08; Oncogene RHO H9; rhoC GTPase; Ras homolog 9 (RhoC); Ras homolog gene family, member A; Ras homolog gene family, member B; Ras homolog gene family, member C; Rho related GTP binding protein RhoC; RHO12; RHOB; RHOC; RHOH9; RP11-426L16.4; Small GTP binding protein RhoC; H9; Rho related GTP binding protein RhoB; ARH12; ARH6; H12.RHOB_HUMAN

研究领域： 细胞生物 信号转导 激酶和磷酸酶

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Chicken, Dog, Rabbit,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需

做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量：22kDa

细胞定位：细胞核 细胞浆 细胞膜

性状：Lyophilized or Liquid

浓度：1mg/ml

免疫原：KLH conjugated synthetic peptide derived from human Rho B:101-196/196

亚型：IgG

纯化方法：affinity purified by Protein A

储存液：0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件：Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PubMed：PubMed

产品介绍： The Rho proteins interact directly with protein kinases which are likely to serve as downstream effector targets of the activated GTPase. The Rho family of GTPases are involved in the regulation of the actin skeleton. Activation of Rho to the GTP bound state promotes formation of stress fibers and is required for the formation and maintenance of focal adhesions. Pathways stimulated by cytokines, hormones and various forms of stress utilize p21 proteins of the Rho family. Possible mediators for these responses include protein kinase such as p120PKN, and p150 ROKa/Rho-A kinase, myosin light chain (MLC) phosphatase and PIP 5-kinase.

Function:

Mediates apoptosis in neoplastically transformed cells after DNA damage. Not essential for development but affects cell adhesion and growth factor signaling in transformed cells. Plays a negative role in tumorigenesis as deletion causes tumor formation. Involved in intracellular protein trafficking of a number of proteins. Targets PKN1 to endosomes and is involved in trafficking of the EGF receptor from late endosomes to lysosomes. Also required for stability and nuclear trafficking of AKT1/AKT which promotes endothelial cell survival during vascular development. Serves as a microtubule-dependent signal that is required for the myosin contractile ring formation during cell cycle cytokinesis. Required for genotoxic stress-induced cell death in breast cancer cells.

Subunit:

Binds ROCK1 and ROCK2. Also binds PKN1/PRK1. Interacts with ARGGEF3, RTKN and AKAP13.

Subcellular Location:

Rho A: Cell membrane; Lipid-anchor; Cytoplasmic side. Cytoplasm; cytoskeleton. Rho B: Late endosome membrane; Lipid-anchor. Cell membrane; Lipid-anchor. Nucleus. Rho C: Cell membrane; Lipid-anchor; Cytoplasmic side.

Post-translational modifications:

Prenylation specifies the subcellular location of RHOB. The farnesylated form is localized to the plasma membrane while the geranylgeranylated form is localized to the endosome.

Similarity:

Belongs to the small GTPase superfamily. Rho family.

SWISS:

P62745

Gene ID:

388

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

产品图片

