

# Rho GTP 酶激活蛋白 32 抗体

产品货号: mIR9296

英文名称: ARHGAP32

中文名称: Rho GTP 酶激活蛋白 32 抗体

别 名: Brain-specific Rho GTPase-activating protein; GAB-associated Cdc42/Rac GTPase-activating protein; GC-GAP; GRIT; GTPase regulator interacting with TrkA; p200RhoGAP; p250GAP; PX-RICS; rac GTPase activating protein; Rho GTPase activating protein 32; Rho-type GTPase-activating protein 32; Rho/Cdc42/Rac GTPase-activating protein RICS; RhoGAP involved in the beta-catenin-N-cadherin and NMDA receptor signaling; RICS.

研究领域: 细胞生物 神经生物学 信号转导 细胞周期蛋白 G蛋白偶联受体 G蛋白信号

抗体来源: Rabbit

克隆类型: Polyclonal

交叉反应: Human, Mouse, Rat, Dog, Pig, Cow, Horse,

产品应用: ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 IF=1:50-200 (石蜡切片需做抗原修复)not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量: 230kDa

细胞定位: 细胞浆 细胞膜

性 状: Lyophilized or Liquid

浓 度: 1mg/ml



免疫原: KLH conjugated synthetic peptide derived from human ARHGAP32:801-920/2078

亚 型: IgG

纯化方法: affinity purified by Protein A

储存液: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件: Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20 °C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PubMed: PubMed

产品介绍: ARHGAP32 is a neuron-associated GTPase-activating protein that may regulate dendritic spine morphology and strength by modulating Rho GTPase.

### **Function:**

GTPase-activating protein (GAP) promoting GTP hydrolysis on RHOA, CDC42 and RAC1 small GTPases. May be involved in the differentiation of neuronal cells during the formation of neurite extensions. Involved in NMDA receptor activity-dependent actin reorganization in dendritic spines. May mediate cross-talks between Ras- and Rho-regulated signaling pathways in cell growth regulation. Isoform 2 has higher GAP activity (By similarity).

## Subunit:

nteracts with NTRK1 (via cytoplasmic domain); the interaction is independent of the phosphorylation state of NTRK1. Interacts with SHC3 (via SH2 domain). Interacts with RASA1 (via SH3 domain); the interaction is necessary for the Ras activation and cell transforming activities of ARHGAP32 (By similarity). Interacts with GAB1 and GAB2. Interacts with CRKL and CRKL. Found in a complex with CRKL and BCAR1; upon EGF stimulation BCAR1 may be replaced by EGFR. Interacts with NCK1 (via SH3 domain); NCK1 recruits phosphorylated BCAR1 to the complex. Isoform 2 interacts with FYN; the interaction appears to be dependent on tyrosine phosphorylation of ARHGAP32. Interacts with EGFR; the interaction requires EGF stimulation and is increased by SHC3. Interacts with CDC42; the interaction requires constitutively active CDC42. Interacts with CTNNB1, DLG4, CDH2 and GRIN2B (By similarity).



## **Subcellular Location:**

Cell junction, synapse, postsynaptic cell membrane, postsynaptic density. Cell projection, dendritic spine By similarity. Cytoplasm, cell cortex. Endosome membrane By similarity. Golgi apparatus membrane By similarity. Endoplasmic reticulum membrane By similarity. Membrane.

## **Tissue Specificity:**

Isoform 1 and isoform 2 are highly expressed in brain and testis. Isoform 1 is also expressed in other tissues such as lung, liver and spleen.

### Post-translational modifications:

Isoform 2 is phosphorylated on multiple tyrosine residues by FYN. Phosphorylated tyrosine residues undergo dephosphorylation after stimulation of NMDA receptors (By similarity). Phosphorylated in vitro by CaMK2 in the presence of calmodulin and calcium; which inhibits GAP activity (By similarity).

## Similarity:

Belongs to the PX domain-containing GAP family.

Contains 1 PX (phox homology) domain.

Contains 1 Rho-GAP domain.

Contains 1 SH3 domain.

**SWISS:** 

A7KAX9

Gene ID:



9743

## Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

# 产品图片

