

## 睾丸凋亡基因抗体

产品货号： mlR1899

英文名称： SPATA17

中文名称： 睾丸凋亡基因抗体

别名： Spermatogenesis-associated protein 17; SPATA17; Spermatogenesis-related protein 11; 1700065F16RIK; 4930504I07Rik; 4930513F16RIK; IQCH; LOC128153; MGC107213; MSRG-11; RGD1565282; RP11-144C20.1; Spata17; SRG11; SPT17\_HUMAN; spermatogenesis associated 17; MSRG11; MSRG-11.

研究领域： 细胞凋亡

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Dog, Cow, Horse,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 IF=1:100-500 （石蜡切片需做抗原修复）  
not yet tested in other applications.  
optimal dilutions/concentrations should be determined by the end user.

分子量： 43kDa

细胞定位： 细胞核

性状： Lyophilized or Liquid

浓度： 1mg/ml

免疫原： KLH conjugated synthetic peptide derived from human SRG11:101-200/361

亚型： IgG

**纯化方法：** affinity purified by Protein A

**储 存 液：** 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

**保存条件：** Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

**PubMed：** PubMed

**产品介绍：** Strongly expressed in adult testis but weakly expressed in the spleen and thymus. Strongly expressed in round and elongating spermatids, and weakly or not expressed in spermatozoa.

**Subcellular Location:**

Cytoplasm.

**Similarity:**

Contains 3 IQ domains.

**SWISS:**

Q96L03

**Gene ID:**

128153

**Important Note:**



This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.