

PRTG 蛋白抗体

产品货号： mlR19447

英文名称： PRTG

中文名称： PRTG 蛋白抗体

别名： FLJ12795; FLJ13221; FLJ25756; IGDC5; Protein Shen-Dan; Protogenin; PRTG; PRTG_HUMAN; RGD1307157.

研究领域： 细胞生物 免疫学 发育生物学 神经生物学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Chicken, Dog, Cow, Horse, Sheep,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 124kDa

细胞定位： 细胞浆 细胞膜

性状： Lyophilized or Liquid

浓度： 1mg/ml

免 疫 原： KLH conjugated synthetic peptide derived from human PRTG:311-410/1150 <Extracellular>

亚 型： IgG

纯化方法： affinity purified by Protein A

储 存 液： 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件： Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed： PubMed

产品介绍： This gene encodes a member of the immunoglobulin superfamily. The encoded transmembrane protein has been associated with the development of various tissues, especially neurogenesis. It has been suggested that this gene may be associated with attention deficit hyperactivity disorder (ADHD). [provided by RefSeq, Nov 2014]

Function:

May play a role in anteroposterior axis elongation

Similarity:

Belongs to the immunoglobulin superfamily. DCC family.

Contains 5 fibronectin type-III domains.

Contains 4 Ig-like (immunoglobulin-like) domains.

SWISS:

Q2VWP7

Gene ID:

283659

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.