

乳腺癌细胞 3 抗体

产品货号： mlR14278

英文名称： DEPDC1B/BRCC3

中文名称： 乳腺癌细胞 3 抗体

别名： BRCC3; Breast cancer cell 3; DEP domain containing 1B; DEP domain containing protein 1B; FLJ11252; DEP1B_HUMAN; HBV X transactivated gene 8 protein; HBV X-transactivated gene 8 protein; HBV XAg-transactivated protein 8; HBxAg transactivated protein 1; OTTHUMP00000122605; OTTHUMP00000221973; XTP1; XTP8.

研究领域： 肿瘤 细胞生物 信号转导 G 蛋白信号

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Cow, Rabbit,

产品应用： WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500

(石蜡切片需做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 62kDa

细胞定位： 细胞浆

性状： Lyophilized or Liquid

浓度： 1mg/ml

免 疫 原 : KLH conjugated synthetic peptide derived from human DEPDC1B/BRCC3:451-529/529

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PubMed : PubMed

产品介绍 : GTPase activator activity. biological process: intracellular signaling cascade

Subcellular Location:

intracellular

Similarity:

Contains 1 DEP domain.

Contains 1 Rho-GAP domain.

SWISS:

Q8WUY9

Gene ID:

55789

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

产品图片

