

过氧化物酶体膜蛋白 4 抗体

产品货号： mlR21115

英文名称： PXMP4

中文名称： 过氧化物酶体膜蛋白 4 抗体

别名： 24 kDa peroxisomal intrinsic membrane protein; Peroxisomal membrane protein 4 (24kD); Peroxisomal membrane protein 4; Peroxisomal membrane protein 4, 24kDa; PMP24; PXMP4_HUMAN.

研究领域： 细胞生物 细胞膜蛋白

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Chicken, Dog, Cow, Zebrafish, Sheep,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量 : 24kDa

细胞定位 : 细胞浆

性 状 : Lyophilized or Liquid

浓 度 : 1mg/ml

免 疫 原 : KLH conjugated synthetic peptide derived from human PXMP4:131-212/212

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : Preservative: 15mM Sodium Azide, Constituents: 1% BSA, 0.01M PBS, pH 7.4

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed : PubMed

产品介绍 background:

PXMP4 belongs to the peroxisomal membrane protein PXMP2/4 family. Hypermethylation-mediated silencing of PXMP4 may be involved in prostate carcinogenesis. PXMP4 undergoes hypermethylation and silencing during the transition of prostate adenocarcinoma cells from androgen dependence to androgen independence.

Subcellular Location:

Peroxisome membrane; Multi pass membrane protein.

SWISS:

Q9Y6I8

Gene ID:

11264

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

产品图片

