

POGK 蛋白抗体

产品货号： mlR12752

英文名称： POGK

中文名称： POGK 蛋白抗体

别 名： BASS 2; BASS2; KIAA1513; KIAA15131; LST003; POGK; POGK_HUMAN; Pogo transposable element with KRAB domain; SLTP003.

研究领域： 细胞生物 免疫学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Dog, Pig, Cow, Horse, Rabbit, Sheep,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量： 69kDa

细胞定位： 细胞核

性 状： Lyophilized or Liquid

浓 度： 1mg/ml

免 疫 原： KLH conjugated synthetic peptide derived from human POGK:461-560/609

亚 型： IgG

纯化方法： affinity purified by Protein A

储 存 液： 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件： Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed： PubMed

产品介绍： The exact function of the protein encoded by this gene is not known. However, this gene product contains a KRAB domain (which is involved in protein-protein interactions) at the N-terminus, and a transposase domain at the C-terminus, suggesting that it may belong to the family of DNA-mediated transposons in human. [provided by RefSeq, Jul 2008]

Subcellular Location:

Nucleus.

Similarity:

Contains 1 DDE domain.

Contains 1 HTH CENPB-type DNA-binding domain.

Contains 1 KRAB domain.

SWISS:

Q9P215

Gene ID:

57645

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.