

血小板衍生生长因子相关蛋白 1 抗体

产品货号： mlR9714

英文名称： PDAP1

中文名称： 血小板衍生生长因子相关蛋白 1 抗体

别名： 28 kDa heat- and acid-stable phosphoprotein; 28kDa heat and acid stable phosphoprotein; HAP28_HUMAN; HASPP28; PAP; PAP1; PDAP 1; Pdap1; PDGF associated protein; PDGF-associated protein; PDGFA associated protein 1; PDGFA-associated protein 1.

研究领域： 细胞生物 信号转导 生长因子和激素

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Rabbit,

产品应用： WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500

（石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 21kDa

细胞定位： 细胞浆 细胞膜

性状： Lyophilized or Liquid

浓度： 1mg/ml

免 疫 原： KLH conjugated synthetic peptide derived from human PDAP1:

亚 型： IgG

纯化方法： affinity purified by Protein A

储 存 液： 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件： Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed： PubMed

产品介绍： The protein encoded by this gene is a phosphoprotein that may upregulate the PDGFA-stimulated growth of fibroblasts and also downregulate the mitogenicity of PDGFB. The encoded protein in rodents has been shown to bind PDGFA with a low affinity. [provided by RefSeq, Dec 2010]

Function:

Enhances PDGFA-stimulated cell growth in fibroblasts, but inhibits the mitogenic effect of PDGFB.

Similarity:

Belongs to the PDAP1 family.

SWISS:

Q13442

Gene ID:

11333

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.