

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

环指蛋白9抗体

产品货号:	mIR9409
英文名称 :	RNF9
中文名称:	环指蛋白 9 抗体
	B30 RING finger protein; Hematopoietic RING finger 1; HERF1; RFB30; RFPL; RING finger protein 9; ite motif containing 10; Tripartite motif protein 10; TRI10_HUMAN.
研究领域:	细胞生物 锌指蛋白
抗体来源:	Rabbit
克隆类型:	Polyclonal
交叉反应 :	Human, Mouse, Rat, Pig, Cow, Sheep, Chimpanzee, mpanzee
产品应用 : 做抗原修复	WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 IF=1:50-200 (石蜡切片需)



分子量: 55kDa

细胞定位: 细胞浆

性 状: Lyophilized or Liquid

浓 度: 1mg/ml

免疫原: KLH conjugated synthetic peptide derived from human TRIM10/RNF9:201-300/481

亚 型: lgG

纯化方法: affinity purified by Protein A

储 存 液: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件: Store at -20 $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20 $^{\circ}$ C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 $^{\circ}$ C.

PubMed: PubMed



产品介绍 background:

Important Note:

TRIM10 is a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. This protein localizes to cytoplasmic bodies. Studies in mice suggest that this protein plays a role in terminal differentiation of erythroid cells.

Function:
Seems to play an important role in erythropoiesis (By similarity).
Subcellular Location:
Cytoplasmic
Similarity:
Belongs to the TRIM/RBCC family.
Contains 1 B box-type zinc finger.
Contains 1 B30.2/SPRY domain.
Contains 1 RING-type zinc finger.
SWISS:
Q9UDY6
Gene ID:
10107



This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.