

## PAPLN 蛋白多糖样硫酸化糖蛋白抗体

产品货号： mIR21079

英文名称： PAPLN

中文名称： PAPLN 蛋白多糖样硫酸化糖蛋白抗体

别名： Papilin; papilin, proteoglycan-like sulfated glycoprotein; PPN\_HUMAN; PRO4977; UNQ2420.

研究领域： 细胞生物 信号转导 泛素

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rabbit,

产品应用： WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500

(石蜡切片需做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 138kDa

细胞定位： 分泌型蛋白

性状： Lyophilized or Liquid

浓度： 1mg/ml

免疫原： KLH conjugated synthetic peptide derived from human PAPLN:1101-1200/1278

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

**PubMed :** PubMed

产品介绍 : PAPLN (Papilin, Proteoglycan Like Sulfated Glycoprotein) is a Protein Coding gene. GO annotations related to this gene include peptidase activity and serine-type endopeptidase inhibitor activity. An important paralog of this gene is ADAMTSL1.

**Subcellular Location:**

Secreted.

**Similarity:**

Belongs to the papilin family.

Contains 1 BPTI/Kunitz inhibitor domain.

Contains 3 Ig-like C2-type (immunoglobulin-like) domains.

Contains 1 PLAC domain.

Contains 5 TSP type-1 domains.

**SWISS:**

O95428

**Gene ID:**

89932

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.