

p53 依赖细胞凋亡调节蛋白抗体

产品货号： mlR17161

英文名称： TP73-AS1

中文名称： p53 依赖细胞凋亡调节蛋白抗体

别名： KIAA0495; p53-dependent apoptosis modulator; PDAM; Putative TP73 antisense gene protein 1; TP73 antisense RNA 1 (non protein coding); TP73 antisense RNA 1. T73AS_HUMAN

研究领域： 细胞生物 细胞凋亡

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量 : 19kDa

细胞定位 : 分泌型蛋白

性 状 : Lyophilized or Liquid

浓 度 : 1mg/ml

免 疫 原 : KLH conjugated synthetic peptide derived from human TP73-AS1:21-120/201

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed : PubMed

产品介绍 May modulate apoptosis via regulation of p53/TP53-dependent anti-apoptotic genes. Induces resistance of the chemotherapy drug cisplatin in glioma cells when underexpressed.

Function:

May modulate apoptosis via regulation of p53/TP53-dependent anti-apoptotic genes. Induces resistance of the chemotherapy drug cisplatin in glioma cells when underexpressed.

Subcellular Location:

Secreted.

SWISS:

Q9UF72

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.