

## 脂肪瘤高迁移率融合蛋白样蛋白 3 抗体

产品货号： mIR18240

英文名称： LHFPL3

中文名称： 脂肪瘤高迁移率融合蛋白样蛋白 3 抗体

别 名： LHFP-like protein 3; lhfp13; LHFPL4; LHPL3\_HUMAN; Lipoma HMGIC fusion partner-like 3 protein; lipoma HMGIC fusion partner-like 4.

研究领域： 细胞生物 免疫学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Dog, Cow, Horse, Sheep,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量： 25kDa

细胞定位： 细胞膜

性 状： Lyophilized or Liquid

浓 度： 1mg/ml

**免 疫 原：** KLH conjugated synthetic peptide derived from human LHFPL3:151-236/236

**亚 型：** IgG

**纯化方法：** affinity purified by Protein A

**储 存 液：** 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

**保存条件：** Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

**PubMed：** PubMed

**产品介绍：** This gene is a member of the lipoma HMGIC fusion partner (LHFP) gene family, which is a subset of the superfamily of tetraspan transmembrane protein encoding genes. Mutations in one LHFP-like gene result in deafness in humans and mice, and a second LHFP-like gene is fused to a high-mobility group gene in a translocation-associated lipoma. A partial gene fragment named LHFPL4 corresponds to a portion of the first exon of this gene. [provided by RefSeq, Jul 2008]

**Subcellular Location:**

Membrane.

**Similarity:**

Belongs to the LHFP family.

**SWISS:**

Q86UP9

**Gene ID:**

375612

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.