

丝氨酸苏氨酸 SRP75 抗体

产品货号： mlR17300

英文名称： SFRS4

中文名称： 丝氨酸/苏氨酸 SRP75 抗体

别名： arginine/serine-rich 4; Pre mRNA splicing factor SRP75; Pre-mRNA-splicing factor SRP75; Serine/arginine-rich splicing factor 4; Splicing factor; Splicing factor, arginine/serine rich 4; SRP001LB; SRP75; Srsf4; SRSF4_HUMAN.

研究领域： 细胞生物 表观遗传学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat,

产品应用： WB=1:500-2000 ELISA=1:500-1000

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 57kDa

细胞定位： 细胞核

性状： Lyophilized or Liquid

浓度： 1mg/ml

免疫原： KLH conjugated synthetic peptide derived from human SFRS4:1-100/494

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PubMed : PubMed

产品介绍 : This gene encodes a member of the arginine/serine-rich splicing factor family. The encoded protein likely functions in mRNA processing. [provided by RefSeq, Feb 2009]

Function:

A probable role in alternative splice site selection during pre-mRNA splicing.

Subcellular Location:

Nucleus speckle.

Post-translational modifications:

Extensively phosphorylated on serine residues in the RS domain.

Similarity:

Belongs to the splicing factor SR family.

Contains 2 RRM (RNA recognition motif) domains.

SWISS:

Q08170

Gene ID:

6429

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

产品图片

