



G 蛋白信号转导调节因子 18 抗体

产品货号 : mlR19858

英文名称 : RGS18

中文名称 : G 蛋白信号转导调节因子 18 抗体

别 名 : Regulator of G protein signalling 13; Regulator of G Protein Signalling 18; Regulator of G-protein signalling 18; RGS 18; RGS13; RGS18; RGS18_HUMAN.

研究领域 : 细胞生物 信号转导 G 蛋白信号

抗体来源 : Rabbit

克隆类型 : Polyclonal

交叉反应 : Human, Mouse, Rat, Dog, Pig, Cow,

产品应用 : WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500
(石蜡切片需做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量 : 28kDa

细胞定位 : 细胞浆

性 状 : Lyophilized or Liquid

浓 度 : 1mg/ml



免 疫 原 : KLH conjugated synthetic peptide derived from human RGS18:1-100/255

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed : PubMed

产品介绍 : This gene encodes a member of the regulator of G-protein signaling family. This protein is contains a conserved, 120 amino acid motif called the RGS domain. The protein attenuates the signaling activity of G-proteins by binding to activated, GTP-bound G alpha subunits and acting as a GTPase activating protein (GAP), increasing the rate of conversion of the GTP to GDP. This hydrolysis allows the G alpha subunits to bind G beta/gamma subunit heterodimers, forming inactive G-protein heterotrimers, thereby terminating the signal. Alternate transcriptional splice variants of this gene have been observed but have not been thoroughly characterized. [provided by RefSeq, Jul 2008]

Function:

Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form. Binds to G(i) alpha-1, G(i) alpha-2, G(i) alpha-3 and G(q) alpha.

Subcellular Location:

Cytoplasm.

Tissue Specificity:

Expressed in peripheral leukocytes, bone marrow, platelet, spleen and fetal liver.



Similarity:

Contains 1 RGS domain.

SWISS:

Q9NS28

Gene ID:

64407

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.