

肿瘤蛋白 p53 诱导蛋白 13(肿瘤抑制基因)

产品货号： mlR6145

英文名称： TP53I13

中文名称： 肿瘤蛋白 p53 诱导蛋白 13(肿瘤抑制基因)

别名： Damage stimulated cytoplasmic protein 1; DSCP1; Tumor protein p53 inducible protein 13; P5I13_HUMAN.

研究领域： 肿瘤 细胞生物 免疫学 细胞周期蛋白

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Dog, Pig, Cow, Horse,

产品应用： WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量 : 39kDa

细胞定位 : 细胞浆 细胞膜

性 状 : Lyophilized or Liquid

浓 度 : 1mg/ml

免 疫 原 : KLH conjugated synthetic peptide derived from human TP53I13:51-150/393

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PubMed : PubMed

产品介绍 background:

TP53I13 may act as a tumor suppressor as it inhibits tumor cell growth, when overexpressed.

Function:

May act as a tumor suppressor. Inhibits tumor cell growth, when overexpressed.

Subcellular Location:

Cell membrane; Single-pass type I membrane protein; Extracellular side (Potential). Cytoplasm. Note=Associates with unknown subcellular structures in the cytoplasm.

Tissue Specificity:

Expressed in heart, placenta, skeletal muscle, testis, brain and lung.

SWISS:

Q8NBRO

Gene ID:

90313

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.