



## 肿瘤/睾丸抗原 2 抗体

产品货号 : mlR6807

英文名称 : CTAG2

中文名称 : 肿瘤/睾丸抗原 2 抗体

别名 : Autoimmunogenic cancer/testis antigen NY ESO 2; Autoimmunogenic cancer/testis antigen NY-ESO-2; CAMEL; Cancer/testis antigen 2; Cancer/testis antigen 6.2; CT 2; CT2; CT6.2; CTAG 2; CTAG2; CTAG2\_HUMAN; CTL recognized antigen on melanoma; ESO 2; ESO2; L antigen family member 1; LAGE 1; LAGE 1a protein; LAGE 2b; LAGE-1; LAGE1; LAGE2b.

研究领域 : 肿瘤 免疫学 肿瘤细胞生物标志物

抗体来源 : Rabbit

克隆类型 : Polyclonal

交叉反应 : Human,

产品应用 : WB=1:500-2000 ELISA=1:500-1000

not yet tested in other applications.



optimal dilutions/concentrations should be determined by the end user.

分子量： 23kDa

细胞定位： 细胞浆

性 状： Lyophilized or Liquid

浓 度： 1mg/ml

免 疫 原： KLH conjugated synthetic peptide derived from human CTAG2:121-210/210

亚 型： IgG

纯化方法： affinity purified by Protein A

储 存 液： 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件： Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PubMed： PubMed



**产品介绍 background:**

Testis and very low level in placenta and in some uterus samples. Observed in 25-50% of tumor samples of melanomas, non-small-cell lung carcinomas, bladder, prostate and head and neck cancers.

**Tissue Specificity:**

Testis and very low level in placenta and in some uterus samples. Observed in 25-50% of tumor samples of melanomas, non-small-cell lung carcinomas, bladder, prostate and head and neck cancers.

**Similarity:**

Belongs to the CTAG family.

**SWISS:**

075638

**Gene ID:**

30848

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

**产品图片**

