



# 肿瘤坏死因子受体相关因子 3 相互作用蛋白 1 抗体

产品货号 : mlR17182

英文名称 : TRAF3IP1

中文名称 : 肿瘤坏死因子受体相关因子 3 相互作用蛋白 1 抗体

别 名 : Interleukin 13 receptor alpha 1 binding protein 1; Interleukin-13 receptor alpha 1-binding protein 1; Microtubule interacting protein that associates with TRAF3; Microtubule-interacting protein associated with TRAF3; MIP T3; MIP-T3; MIPT3; MIPT3\_HUMAN; TNF receptor associated factor 3 interacting protein 1; TRAF3-interacting protein 1; Traf3ip1.

研究领域 : 细胞生物 免疫学 信号转导

抗体来源 : Rabbit

克隆类型 : Polyclonal

交叉反应 : Human, Mouse, Rat, Chicken, Dog, Pig, Cow, Horse, Rabbit, Zebrafish, Sheep, Guinea Pig,



**产品应用** : ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 (石蜡切片需做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

**分子量** : 79kDa

**细胞定位** : 细胞浆

**性 状** : Lyophilized or Liquid

**浓 度** : 1mg/ml

**免 疫 原** : KLH conjugated synthetic peptide derived from human TRAF3IP1:501-600/691

**亚 型** : IgG

**纯化方法** : affinity purified by Protein A

**储 存 液** : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

**保存条件** : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When



reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

**PubMed :** PubMed