

肿瘤坏死因子受体相互作用蛋白抗体

产品货号： mIR4983

英文名称： TRIP

中文名称： 肿瘤坏死因子受体相互作用蛋白抗体

别名： RING finger protein 206; RNF206; TRAF Interacting Protein; TRAIP; TRIP.

研究领域： 肿瘤 转录调节因子 细胞膜受体

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Dog, Pig, Horse,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 53kDa

细胞定位： 细胞膜

性状： Lyophilized or Liquid

浓度： 1mg/ml

免疫原： KLH conjugated synthetic peptide derived from human TRIP:201-300/469

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed : PubMed

产品介绍 : RING finger motif and a putative coiled-coil domain. A similar murine protein interacts with TNFR-associated factor 1 (TRAF1), TNFR-associated factor 2 (TRAF2), and cylindromatosis. The interaction with TRAF2 inhibits TRAF2-mediated nuclear factor kappa-B, subunit 1 activation that is required for cell activation and protection against apoptosis.

Function:

Inhibits activation of NF-kappa-B mediated by TNF.

Subunit:

Binds TRAF1 and TRAF2 and is part of the receptor-TRAF signaling complex. Interacts with CYLD.

Subcellular Location:

Cytoplasm. Cytoplasm, perinuclear region.

Similarity:

Contains 1 RING-type zinc finger.

SWISS:

Q9BWF2

Gene ID:

10293

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.