

not yet tested in other applications.

FBXL17 蛋白抗体

产品货号: mlR8469	
英文名称 : FBXL17	
中文名称: FBXL17 蛋白抗体	
别 名: F box and leucine rich repeat protein 17; F box only p and leucine-rich repeat protein 17; F-box only protein 13; F-box/LFBXO13; FBXL13; FXL17_HUMAN; DKFZP434C1715.	
研究领域: 细胞生物 免疫学 泛素	
抗体来源: Rabbit	
克隆类型: Polyclonal	
交叉反应 : Human, Mouse, Rat, Chicken, Dog, Pig, Cow, Horse, l	Rabbit, Sheep,
产品应用: WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 做抗原修复)	IHC-F=1:400-800 IF=1:50-200 (石蜡切片需



optimal dilutions/concentrations should be determined by the end user.

分子量: 76kDa

细胞定位: 细胞核

性 状: Lyophilized or Liquid

浓 度: 1mg/ml

免疫原: KLH conjugated synthetic peptide derived from human FBXL17:451-550/701

亚 型: lgG

纯化方法: affinity purified by Protein A

储存液: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件: Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20 °C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PubMed: PubMed



产品介绍: Members of the F-box protein family, such as FBXL17, are characterized by an approximately 40-amino acid F-box motif. SCF complexes, formed by SKP1 (MIM 601434), cullin (see CUL1; MIM 603134), and F-box proteins, act as protein-ubiquitin ligases. F-box proteins interact with SKP1 through the F box, and they interact with ubiquitination targets through other protein interaction domains (Jin et al., 2004 [PubMed 15520277]).[supplied by OMIM, Mar 2008]

Function:
Substrate-recognition component of the SCF (SKP1-CUL1-F-box protein)-type E3 ubiquitin ligase complex.
Subunit:
Interacts with SKP1 and CUL1
Similarity:
Contains 1 F-box domain.
SWISS:
Q9UF56
Gene ID:
64839

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.



产品图片

