

肝癌易感蛋白 3 抗体

产品货号： mlR19578

英文名称： PSMG2/HCCA3

中文名称： 肝癌易感蛋白 3 抗体

别名： CD40 ligand activated specific transcript 3; CLAST3; HCCA3; Hepatocellular carcinoma susceptibility protein; Hepatocellular carcinoma-susceptibility protein 3; HSPC260; HsT1707; Likely ortholog of mouse CD40 ligand activated specific transcript 3; MDS003; PAC-2; PAC2; proteasome (prosome, macropain) assembly chaperone 2; Proteasome assembly chaperone 2; psmG2; PSMG2_HUMAN; TNFSF5IP1; Tumor necrosis factor superfamily member 5-induced protein 1.

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Cow,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 29kDa

细胞定位： 细胞核

性状： Lyophilized or Liquid

浓度： 1mg/ml

免疫原： KLH conjugated synthetic peptide derived from human PSMG2/HCCA3:101-200/264

亚型： IgG

纯化方法： affinity purified by Protein A

储存液： 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件： Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed： PubMed