

L-丝氨酸脱氨酶抗体

产品货号： mIR19607

英文名称： SDSL

中文名称： L-丝氨酸脱氨酶抗体

别 名： L serine deaminase; L serine dehydratase/L threonine deaminase; L threonine dehydratase; L-serine deaminase; L-serine dehydratase/L-threonine deaminase; L-threonine dehydratase; SDH 2; SDH; SDH1; SDS RS1; Sdsl; SDSL_HUMAN; Serine dehydratase 2; Serine dehydratase like; Serine dehydratase.

研究领域： 细胞生物 信号转导 干细胞

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Dog, Pig, Cow,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量： 35kDa

细胞定位： 细胞浆

性 状： Lyophilized or Liquid

浓 度： 1mg/ml

免 疫 原： KLH conjugated synthetic peptide derived from human SDSL:61-160/329

亚 型： IgG

纯化方法： affinity purified by Protein A

储 存 液： 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件： Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed： PubMed

产品介绍： Has low serine dehydratase and threonine dehydratase activity.

Function:

Has low serine dehydratase and threonine dehydratase activity.

Subunit:

Homodimer.

Similarity:

Belongs to the serine/threonine dehydratase family.

SWISS:

Q96GA7

Gene ID:

113675

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.