

## LYRM2 蛋白抗体

产品货号： mlR18583

英文名称： LYRM2

中文名称： LYRM2 蛋白抗体

别 名： 2610208E05Rik; DJ12208.2; FLJ18193; FLJ18317; FLJ21130; LYR motif-containing protein 2; lyrm2;  
LYRM2\_HUMAN; MGC104253; RP23-301P13.1.

研究领域： 细胞生物 免疫学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Dog, Cow, Sheep,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需  
做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量： 10kDa

细胞定位： 细胞核

性 状： Lyophilized or Liquid

浓 度： 1mg/ml

**免 疫 原：** KLH conjugated synthetic peptide derived from human LYRM2:21-80/88

**亚 型：** IgG

**纯化方法：** affinity purified by Protein A

**储 存 液：** 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

**保存条件：** Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

**PubMed：** PubMed

**产品介绍：** LYRM2 is an 88 amino acid protein belonging to the complex I LYR family. LYRM2 is encoded by a gene located on human chromosome 6, which contains 170 million base pairs and comprises nearly 6% of the human genome. Deletion of a portion of the q arm of chromosome 6 is associated with early onset intestinal cancer, suggesting the presence of a cancer susceptibility locus. Additionally, Porphyria cutanea tarda, Parkinson's disease, Stickler syndrome and a susceptibility to bipolar disorder are all associated with genes that map to chromosome 6.

**Similarity:**

Belongs to the complex I LYR family.

**SWISS:**

Q9NU23

**Gene ID:**

57226



**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.