

KIN17 抗体

产品货号： mIR17062

英文名称： KIN17

中文名称： KIN17 抗体

别名： antigenic determinant of recA protein homolog; Binding to curved DNA; BTCD; DNA/RNA-binding protein KIN17; HsKin17 protein; KIN; KIN antigenic determinant of recA protein homolog mouse; Kin17 protein; KIN17_HUMAN.

研究领域： 细胞生物 细胞周期蛋白 表观遗传学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Dog, Cow, Horse, Sheep,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需

做抗原修复)

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量 : 45kDa

细胞定位 : 细胞核 细胞浆

性 状 : Lyophilized or Liquid

浓 度 : 1mg/ml

免 疫 原 : KLH conjugated synthetic peptide derived from human KIN17:301-393/393

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed : PubMed

产品介绍 The protein encoded by this gene is a nuclear protein that forms intranuclear foci during proliferation and is redistributed in the nucleoplasm during the cell cycle. Short-wave ultraviolet light provokes the relocalization of the protein, suggesting its participation in the cellular response to DNA damage. Originally selected based on protein-binding with RecA antibodies, the mouse protein presents a limited similarity with a functional domain of the bacterial RecA protein, a characteristic shared by this human ortholog. Alternative splicing of this gene results in multiple transcript variants. [provided by RefSeq, Jan 2012]

Function:

Involved in DNA replication and the cellular response to DNA damage. May participate in DNA replication factories and create a bridge between DNA replication and repair mediated by high molecular weight complexes. May play a role in illegitimate recombination and regulation of gene expression. May participate in mRNA processing. Binds, in vitro, to double-stranded DNA. Also shown to bind preferentially to curved DNA in vitro and in vivo (By similarity). Binds via its C-terminal domain to RNA in vitro.

Subcellular Location:

Nucleus. Cytoplasm. During S phase, strongly associated with the nuclear matrix, and to chromosomal DNA in the presence of DNA damage. Also shows cytoplasmic localization in elongated spermatids.

Tissue Specificity:

Ubiquitously expressed in all tissues examined, with highest levels in skeletal muscle, heart and testis. Differentially expressed in non-tumorigenic and tumorigenic cell lines. Highly expressed in proliferating epithelial keratinocyte cells in vitro (at protein level).

Similarity:

Belongs to the KIN17 family.

Contains 1 C2H2-type zinc finger.

SWISS:

O60870

Gene ID:

22944

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.