



KIAA1751 蛋白抗体

产品货号 : mlR17027

英文名称 : KIAA1751

中文名称 : KIAA1751 蛋白抗体

别 名 : FLJ45476; Hypothetical protein LOC85452; K1751_HUMAN; KIAA1751; RP1-140A9.6;
Uncharacterized protein KIAA1751.

研究领域 : 细胞生物 免疫学

抗体来源 : Rabbit

克隆类型 : Polyclonal

交叉反应 : Human, Mouse, Rat,

产品应用 : ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 (石蜡切片需
做抗原修复)



not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分子量： 87kDa

细胞定位： 细胞浆

性 状： Lyophilized or Liquid

浓 度： 1mg/ml

免 疫 原： KLH conjugated synthetic peptide derived from human KIAA1751:181-280/762

亚 型： IgG

纯化方法： affinity purified by Protein A

储 存 液： 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件： Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.



PubMed : PubMed

产品介绍 : KIAA1751 is a 762 amino acid uncharacterized protein. Existing as three alternatively spliced isoforms, KIAA1751 is encoded by a gene that maps to human chromosome 1p36.33. Chromosome 1 spans 260 million base pairs, contains over 3,000 genes and comprises nearly 8% of the human genome. Chromosome 1 houses a large number of disease-associated genes, including those that are involved in familial adenomatous polyposis, Stickler syndrome, Parkinson's disease, Gaucher disease, schizophrenia and Usher syndrome. Aberrations in chromosome 1 are found in a variety of cancers, including head and neck cancer, malignant melanoma and multiple myeloma.

SWISS:

Q9C0B2

Gene ID:

85452

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.