

## KIAA1429 蛋白抗体

产品货号： mlR17003

英文名称： KIAA1429

中文名称： KIAA1429 蛋白抗体

别名： DKFZp434I116; DKFZp781B2117; fSAP121; KIAA1429; MGC138493; MGC141940; MSTP054; Protein virilizer homolog; VIR\_HUMAN.

研究领域： 细胞生物 信号转导 转录调节因子 表观遗传学

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Cow, Horse, Rabbit, Sheep, Rhesus monkey, Gorilla, Orangutan, Platypus

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量 : 202kDa

细胞定位 : 细胞核

性 状 : Lyophilized or Liquid

浓 度 : 1mg/ml

免 疫 原 : KLH conjugated synthetic peptide derived from human KIAA1429:1661-1760/1812

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

**PubMed :** PubMed

**产品介绍 :** Made up of nearly 146 million bases, chromosome 8 encodes about 800 genes. Translocation of portions of chromosome 8 with amplifications of the c-Myc gene are found in some leukemias and lymphomas, and typically associated with a poor prognosis. Portions of chromosome 8 have been linked to schizophrenia and bipolar disorder. Trisomy 8, also known as Warkany syndrome 2, most often results in early miscarriage but is occasionally seen in a mosaic form in surviving patients who suffer to a varying degree from a number of symptoms including retarded mental and motor development, and certain facial and developmental defects. WRN is a DNA helicase encoded by chromosome 8 and shown defective in those with the early aging disorder Werner syndrome. Chromosome 8 is also associated with Pfeiffer syndrome, congenital hypothyroidism and Waardenburg syndrome. The KIAA1429 gene product has been provisionally designated KIAA1429 pending further characterization.

**Function:**

May be involved in mRNA splicing regulation.

**Subcellular Location:**

Nucleus.

**Similarity:**

Belongs to the vir family.

**SWISS:**

Q69YN4

**Gene ID:**

25962

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.