



KIAA1109 蛋白抗体

产品货号 : mlR16990

英文名称 : KIAA1109

中文名称 : KIAA1109 蛋白抗体

别 名 : Fragile site-associated protein; K1109_HUMAN; Kiaa1109; Uncharacterized protein KIAA1109.

研究领域 : 肿瘤 细胞生物 免疫学

抗体来源 : Rabbit

克隆类型 : Polyclonal

交叉反应 : Human, Mouse, Rat, Cow, Rabbit,

产品应用 : ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 (石蜡切片需做抗原修复)

not yet tested in other applications.



optimal dilutions/concentrations should be determined by the end user.

分子量： 555kDa

细胞定位： 细胞核

性 状： Lyophilized or Liquid

浓 度： 1mg/ml

免 疫 原： KLH conjugated synthetic peptide derived from human KIAA1109:4901-5005//5005

亚 型： IgG

纯化方法： affinity purified by Protein A

储 存 液： 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件： Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

PubMed： PubMed



产品介绍 : This gene is located on the long arm of chromosome 4 in a region that is associated with susceptibility to celiac disease. The encoded protein is similar to a Chinese hamster protein that is associated with spermatocyte and adipocyte differentiation. The C-terminus of the protein is also similar to a *Caenorhabditis elegans* protein that plays a role in lipid storage. In mammals, this protein is thought to function in the regulation of epithelial growth and differentiation, and in tumor development. [provided by RefSeq, Oct 2009]

Subcellular Location:

Membrane.

Tissue Specificity:

Highly expressed in testis and ovary. Weakly or not expressed in other tissues.

SWISS:

Q2LD37

Gene ID:

84162

Important Note:

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.