

## 粘蛋白 19 抗体

产品货号： mIR17905

英文名称： MUC19

中文名称： 粘蛋白 19 抗体

别 名： Gel-forming secreted mucin-19; MUC-19; Muc19; MUC19\_MOUSE; mucin apoprotein; Mucin-19; Sublingual apomucin.

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Mouse, Rat,

产品应用： WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500

（石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量： 690kDa

细胞定位： 分泌型蛋白

性 状： Lyophilized or Liquid

浓 度： 1mg/ml

免 疫 原： KLH conjugated synthetic peptide derived from mouse MUC19:7401-7524/7524

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

**PubMed :** PubMed

产品介绍 : May function in ocular mucus homeostasis.

**Subcellular Location:**

Secreted.

**Tissue Specificity:**

Specifically expressed in sublingual salivary glands. Expressed by mucous cells of the submandibular gland and submucosal gland of the trachea. Expression is altered in sld (sublingual gland differentiation arrest) mutants.

**Similarity:**

Contains 1 CTCK (C-terminal cystine knot-like) domain.

Contains 1 TIL (trypsin inhibitory-like) domain.

Contains 2 VWFC domains.

Contains 3 VWFD domains.

**Gene ID:**

239611

**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.