

## Kelch 样蛋白 4 抗体

产品货号： mlR16772

英文名称： KLHL4

中文名称： Kelch 样蛋白 4 抗体

别名： DKELCHL; Kelch (Drosophila) like 4; Kelch-like protein 4; KHL4; KLHL4; KLHL4\_HUMAN.

研究领域： 细胞生物 信号转导 结合蛋白 细胞骨架

抗体来源： Rabbit

克隆类型： Polyclonal

交叉反应： Human, Mouse, Rat, Pig, Cow, Horse, Rabbit, Sheep,

产品应用： ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 （石蜡切片需做抗原修复）

not yet tested in other applications.

optimal dilutions/concentrations should be determined by the end user.

分 子 量 : 80kDa

细胞定位 : 细胞浆

性 状 : Lyophilized or Liquid

浓 度 : 1mg/ml

免 疫 原 : KLH conjugated synthetic peptide derived from human KLHL4:601-700/718

亚 型 : IgG

纯化方法 : affinity purified by Protein A

储 存 液 : 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件 : Store at -20 ° C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20° C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 ° C.

**PubMed :** PubMed

**产品介绍 background:**

This gene encodes a member of the kelch family of proteins, which are characterized by kelch repeat motifs and a POZ/BTB protein-binding domain. It is thought that kelch repeats are actin binding domains. However, the specific function of this protein has not been determined. Alternative splicing of this gene results in two transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008]

**Subcellular Location:**

Cytoplasm > cytoskeleton.

**Tissue Specificity:**

Expressed in adult fibroblasts and in a range of fetal tissues including tongue, palate, and mandible.

**Similarity:**

Contains 1 BTB (POZ) domain.

Contains 6 Kelch repeats.

**SWISS:**

Q9C0H6

**Gene ID:**

56062



**Important Note:**

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.