

KCTD6 蛋白抗体

产品货号: mIR16936

英文名称: KCTD6

中文名称: KCTD6 蛋白抗体

别 名: BTB/POZ domain-containing protein KCTD6; kctd6; KCTD6_HUMAN.

研究领域: 细胞生物 免疫学 神经生物学 通道蛋白

抗体来源: Rabbit

克隆类型: Polyclonal

交叉反应: Human, Mouse, Rat, Pig, Cow, Sheep,

产品应用: WB=1:500-2000 ELISA=1:500-1000 IHC-P=1:400-800 IHC-F=1:400-800 ICC=1:100-500 IF=1:100-500 (石蜡切片需做抗原修复)

not yet tested in other applications.



optimal dilutions/concentrations should be determined by the end user.

分子量: 28kDa

细胞定位: 细胞膜

性 状: Lyophilized or Liquid

浓 度: 1mg/ml

免疫原: KLH conjugated synthetic peptide derived from human KCTD6:101-200/237

亚 型: IgG

纯化方法: affinity purified by Protein A

储存液: 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

保存条件: Store at -20 $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20 $^{\circ}$ C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 $^{\circ}$ C.

PubMed: PubMed



applications.

产品介绍 KCTD6 (potassium channel tetramerisation domain containing 6) is a 237 amino acid protein that contains one BTB domain, suggesting a possible role as a transcriptional regulator. The gene encoding KCTD6 maps to chromosome 3, which contains over 1,100 genes. Notably, a chemokine receptor gene cluster and a variety of human cancer related loci reside on chromosome 3. Particular regions of the chromosome 3 short arm are deleted in many types of cancer cells as well.

Similarity:	
Contains 1 BTB (POZ) domain.	
swiss:	
Q8NC69	
Gene ID:	
200845	
mportant Note:	

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic